



## Comments and Controversies

## Do we need to revise the tripartite subdivision hypothesis of the human subthalamic nucleus (STN)?



Anneke Alkemade, Birte U. Forstmann\*

Cognitive Science Center Amsterdam, University of Amsterdam, Netherlands

## ARTICLE INFO

## Article history:

Accepted 7 March 2014

Available online 15 March 2014

## Keywords:

Subthalamic nucleus

Basal ganglia

7 T magnetic resonance imaging

Deep brain stimulation

## ABSTRACT

The exciting development of ultra-high resolution 7 Tesla (T) magnetic resonance imaging (MRI) has made it possible to clearly visualize and delineate the subthalamic nucleus (STN). Ultra-high resolution MRI provides a first step in the ongoing improvement of imaging techniques rendering it likely that in the near future specific subareas of small brain nuclei such as the STN can be visualized. These developments can contribute to improve clinical imaging, allowing even more accurate targeting of the STN. This is interesting in view of putative limbic, associative, and sensorimotor subdivisions within the STN. The concept of anatomically distinct subdivisions is attractive, both from an anatomical as well as a clinical perspective. However, we argue that the current leading hypothesis of three STN subdivisions is based on low numbers of clinical observations and primate tracing studies. 7 T imaging provides us with markers that could potentially help us to distinguish subdivisions, but our preliminary findings do not indicate the existence of subdivisions. In our opinion additional research is needed. As a consequence the tripartite hypothesis should therefore still be a topic of debate. In view of the possible clinical implications, we would like to raise the question whether anatomical evidence on the topological organization within the STN points towards delineated subdivisions, or an organization without strict anatomical boundaries or septa. The latter would require a revision of the current tripartite hypothesis of the human STN.

© 2014 Elsevier Inc. All rights reserved.

The STN is a small but important brain structure within the basal ganglia. It plays a crucial role in sensorimotor, cognitive, and limbic functions (Temel et al., 2005). The STN serves as a surgical target for deep brain stimulation (DBS) to provide symptomatic relief in a number of movement disorders refractory to pharmacological intervention (Rughani et al., 2013). Interestingly, DBS of the STN in addition to its clear effects on sensorimotor functions also can cause apathy, compulsivity, hypersexuality, cognitive dysfunction, as well as clinical depression including suicidal ideation (Temel et al., 2005). These effects have been ascribed to the existence of putatively anatomically distinct, sensorimotor, cognitive, and limbic subparts of the human STN (Parent and Hazrati, 1995a, b), although an alternative explanation for the effects of DBS may be the interferences with serotonergic signaling by high frequency stimulation (Navailles et al., 2010; Temel et al., 2007). Additionally, we cannot exclude that cell groups and fibers outside the STN may be affected (Munding, 1965). According to the tripartite STN division, the dorsolateral part of the nucleus is responsible for sensorimotor functions, the central part of the STN fulfills cognitive functions, and the medial tip is involved in limbic functions. The existence of distinct STN subdivisions forms the current leading hypothesis

in STN neuroanatomy, and is attractive from a conceptual point of view. Theoretically unwanted side-effects of STN DBS could be minimized or even prevented by specific targeting of the sensorimotor subdivision of the STN. Although in today's clinical practice, the electric current field is most likely to extend the borders of the STN. One could speculate that with further technical developments of STN imaging techniques, or microelectrode recordings during surgery in patients it will become possible to identify and target these putative STN subdivisions in humans.

However, the literature indicates that the underlying neuroanatomy is more complicated, and it is feasible that the hypothesis of the tripartite STN division represents an oversimplification of the anatomical make-up (for review see (Keuken et al., 2012)). We therefore asked the following questions: Is there conclusive evidence supporting the tripartite organization of subdivisions within the human STN? Or is there a need to revise current hypotheses on the topological organization within the human STN?

Support for the tripartite organization of the STN in humans is largely based on the extrapolation of studies in nonhuman primates, on a number of clinical observations, and a recent study using diffusion weighted imaging in humans (Haynes and Haber, 2013; Lambert et al., 2012; Parent, 1990; Temel et al., 2005). Primate studies using tracings to map STN projections have formed the basis for the tripartite hypothesis of the STN. These results of these studies have been interpreted

\* Corresponding author at: Cognitive Science Center Amsterdam, University of Amsterdam, Nieuwe Achtergracht 129, 1018WS Amsterdam, Netherlands.

E-mail address: [bforstmann@gmail.com](mailto:bforstmann@gmail.com) (B.U. Forstmann).

by others in well-cited reviews, contributing to the acceptance of the tripartite subdivision theory (Alexander and Crutcher, 1990; Haynes and Haber, 2013; Joel and Weiner, 1997; Parent, 1990; Parent and Hazrati, 1995a, b). Although extremely valuable, tracing studies are technically challenging and have a number of inherent technical limitations. In addition, large discrepancies exist between individual tracing studies (Keuken et al., 2012). Findings are strongly dependent on tracer injection volume and size of the injection site. Since injection sites are generally small to prevent leakage to adjacent brain structures, anatomical connections may be only partly uncovered, and the size of projection sites may therefore be underestimated. In addition, the number of reported observations in primates is usually limited. A recent, and very extensive study by Haynes and Haber (Haynes and Haber, 2013), provides crucial anatomical detail on the primate STN. This study supports the existence of a topological organization within the STN, although strong overlap between the three proposed functional areas exists (Haynes and Haber, 2013). It should be noted that even though the study by Haynes and Haber is of high quality, it is again based on a limited number (2–4) of observations per injection site, and it remains difficult to determine to what extent observations from nonhuman primates can be extrapolated to the human situation.


One may argue that a lack of anatomical organization does not exclude a functional organization. Clinical observations have shown that a hypomanic state in two patients was caused by stimulation via the DBS contact located in the anteromedial STN. The anteromedial contact, as well as a contact more dorsal improved motor symptoms. Contacts at the boundaries of the STN affected neither behavior, nor motor performance (Mallet et al., 2007). These clinical descriptions argue for topological differences, even though these studies are complicated by variations in electrode positioning (Okun et al., 2005), and variations in the orientation of the electrode or STN axis (Plaha et al., 2006). Apart from clinical observations, to our knowledge, only one study has addressed the tripartite division within the human STN. Lambert and colleagues report conformation for three functional zones in the STN using diffusion weighted imaging (DWI). However, this evidence was only obtained for 14 out of 24 STN's tested in these studies ((Lambert et al., 2012) supplements). Interestingly, a substantial number of STN voxels in all tested STN's met the criteria of more than one functional subdivision. The authors did not provide any information on the spatial location of these specific voxels, but in view of the large numbers of voxels meeting the criteria for multiple subdivisions based on their regional connectivity, it is unlikely that these voxels are exclusively located exactly at the interface between two functional areas, and that these voxels therefore contain connectivity patterns meeting the criteria for more than one subdivision. Although these DWI-studies, for the first

time, investigate connectivity profiles in the human STN of healthy subjects, the data shows large variation. In 10 out of 24 STN's there is no evidence for three functional zones within the STN. We feel that the data reported by Lambert et al. may, to some extent, support a topological organization within the STN, but there is no strong support for a tripartite organization.

An alternative approach would be to investigate evidence from the rodent STN, as a model for the human STN. However, data from rodents are difficult to interpret since the STN in rats is an open nucleus with dendrites extending into brain areas outside the STN (Afsharpoor, 1985a, b). Contrary to rodents, in primates, including humans, the nucleus is considered to be closed with the dendrites largely restricted to the STN (Rafols and Fox, 1976). In addition to the interspecies differences within the STN, the input and output areas of the STN may also show significant interspecies differences. A recent review of animal models on Parkinson's disease indicates that at present no model is available that faithfully reproduces all features of human Parkinson's disease (Lee et al., 2012). In view of these interspecies differences we are not confident that rodent models will provide us with the answer to our question whether there is a tripartite organization of subdivisions within the human STN.

Although many valuable studies have been published addressing the question whether STN subdivisions are present, the need for further studies on the human STN becomes clear. In addition, the following questions arise: Can the human STN and its putative subdivisions be visualized using techniques available for research and clinical practice within the near future? Will it be feasible to test the hypothesis of a human tripartite STN organization?

With the advent of ultra-high resolution 7 Tesla (T) magnetic resonance imaging (MRI) techniques, it is now possible to visualize the human STN in vivo for research purposes (Cho et al., 2010; de Hollander et al., in press; Forstmann et al., 2010, 2012; Keuken et al., 2013; Massey et al., 2012) (see Fig. 1). However, it will take time before these state-of-the-art imaging techniques can be applied routinely for research as well as diagnostics and surgical planning since 7 T MRI scanners are not widely available. In addition, more research is warranted on MR chemistry. Interesting recent publications indicate that using quantitative susceptibility mapping (QSM) have made it possible to quantify brain iron content (Deistung et al., 2013; Langkammer et al., 2012; Schweser et al., 2011). Other methods such as microelectrode recordings in patients during surgery for identification of subdivisions are also promising (Zaidel et al., 2010). Tracing studies as performed in non-human primates cannot be performed in humans for obvious reasons. Recent developments in optogenetic techniques will provide more insight in the molecular events occurring in the STN and are likely to



**Fig. 1.** Illustration of STN visualization using T2\*-weighted 0.5 mm<sup>3</sup> isotropic 7 T MRI. A: Transverse view; B: Coronal view; and C: Sagittal view.

contribute to the improvement of the general understanding of the STN (Gradinaru et al., 2009), but are restricted to animal models.

So far, the highest level of detail on the functional neuroanatomy of the STN has been obtained by histological and immunocytochemical investigations. Since these types of studies are scarce, we report both studies on human as well as nonhuman primates. We would like to note that these anatomical studies are usually descriptive, and studies are generally more focused on the dorsolateral, central and medial aspects of the nucleus, rather than the antero-posterior axis. The STN is considered to be a mainly glutamatergic nucleus (Smith et al., 1990). Immunocytochemical glutamatergic markers in monkeys (Kuwajima et al., 2004), enkephalin markers in monkeys (Aubert et al., 2007), GABA-ergic markers in monkeys and humans (Augood et al., 1999; Charara et al., 2000; Kultas-Ilinsky et al., 1998; Levesque and Parent, 2005), dopaminergic markers in humans (Augood et al., 2000; Hedreen, 1999; Hurd et al., 2001) do not show a clear tripartite organization within the STN. Serotonergic markers in humans and monkeys (Mori et al., 1985; Parent et al., 2011), as well as expression of the calcium binding proteins parvalbumin and calretinin in humans (Augood et al., 1999; Parent et al., 1996) show an inhomogeneous expression in the STN. Immunoreactivity has not been coupled to any specific function within the STN. Markers were distributed over the entire nucleus. However, parvalbumin staining was more prominent in the dorsolateral part of the STN, whereas calretinin revealed a higher expression pattern in the ventromedial part (Augood et al., 1999; Parent et al., 1996). Serotonin was mainly present in the ventral and the medial part of the nucleus, and the serotonin transporter was more intense in the anterior part of the STN. The description of the distribution of serotonin and its transporter may reflect a mismatch in their location, but is more likely to reflect a difference in the interpretation of the experimental data by the individual research groups. In conclusion, a number of neuronal markers show increased concentrations of immunoreactivity in specific subparts of the STN. This supports a topological organization without strict anatomical boundaries within the nucleus, rather than the existence of three functionally and anatomically distinct subdivisions. One should, however, keep in mind that we cannot exclude that different markers for the STN will show a more distinct distribution within the STN.

The recent study by Lambert et al. (2012), is, to our knowledge, the only study showing in vivo imaging of anatomically distinct subdivisions within the STN. We, however, have not been able to find support for STN subdivisions in our own studies ((de Hollander); Alkemade and Forstmann unpublished data). It is feasible that due to anatomical overlap in the functional organization within the STN, it will remain impossible to distinguish STN subdivisions. In addition, the anatomical overlap hampers selective targeting of the motor subdivision of the STN in DBS surgery.

Some neurosurgical teams report a level of technical precision allowing them to specifically target the dorsolateral part of the STN (Greenhouse et al., 2011; Mallet et al., 2007). If a tripartite organization of subdivisions within the STN exists, this would allow more exclusive targeting of the sensorimotor part of the STN. Does this surgical precision provide such therapeutic benefit? In our opinion, although conceptually attractive there is not enough evidence to support the tripartite subdivision of the human STN. We therefore argue that the tripartite hypothesis should be revised. At present, evidence points towards a topological organization within the STN without strict anatomical borders or septa. We propose that there is at least partial overlap between putative functional zones within the STN. At present it is unclear to what extent functional zones overlap, but it is feasible that the overlap is substantial which has clear implications for basic science as well as for DBS of the STN.

## Acknowledgments

The authors would like to thank Max C Keuken for valuable discussions. This research line is financially supported by the European

Research Council (313481) (BUF), and the Dutch Brain Foundation (AA and BUF).

## References

- Afsharpoor, S., 1985a. Light microscopic analysis of Golgi-impregnated rat subthalamic neurons. *J. Comp. Neurol.* 236, 1–13.
- Afsharpoor, S., 1985b. Topographical projections of the cerebral cortex to the subthalamic nucleus. *J. Comp. Neurol.* 236, 14–28.
- Alexander, G.E., Crutcher, M.D., 1990. Functional architecture of basal ganglia circuits: neural substrates of parallel processing. *Trends Neurosci.* 13, 266–271.
- Aubert, I., Guigoni, C., Li, Q., Dovero, S., Bioulac, B.H., Gross, C.E., Crossman, A.R., Bloch, B., Bezard, E., 2007. Enhanced preproenkephalin-B-derived opioid transmission in striatum and subthalamic nucleus converges upon globus pallidus internalis in L-3,4-dihydroxyphenylalanine-induced dyskinesia. *Biol. Psychiatry* 61, 836–844.
- Augood, S.J., Waldvogel, H.J., Munkle, M.C., Faull, R.L., Emson, P.C., 1999. Localization of calcium-binding proteins and GABA transporter (GAT-1) messenger RNA in the human subthalamic nucleus. *Neuroscience* 88, 521–534.
- Augood, S.J., Hollingsworth, Z.R., Standaert, D.G., Emson, P.C., Penney Jr., J.B., 2000. Localization of dopaminergic markers in the human subthalamic nucleus. *J. Comp. Neurol.* 421, 247–255.
- Charara, A., Heilman, T.C., Levey, A.I., Smith, Y., 2000. Pre- and postsynaptic localization of GABA(B) receptors in the basal ganglia in monkeys. *Neuroscience* 95, 127–140.
- Cho, Z.H., Min, H.K., Oh, S.H., Han, J.Y., Park, C.W., Chi, J.G., Kim, Y.B., Paek, S.H., Lozano, A.M., Lee, K.H., 2010. Direct visualization of deep brain stimulation targets in Parkinson disease with the use of 7-tesla magnetic resonance imaging. *J. Neurosurg.* 113, 639–647.
- de Hollander, G., Keuken, M.C., Bazin, P.L., Weiss, M., Neumann, J., Reimann, K., Wähnert, M., Turner, R., Forstmann, B.U., Schäfer, A., 2014. A gradual increase of iron towards the medial-inferior tip of the subthalamic nucleus. *Hum. Brain Mapp.* (in press).
- Deistung, A., Schafer, A., Schweser, F., Biedermann, U., Turner, R., Reichenbach, J.R., 2013. Toward in vivo histology: a comparison of quantitative susceptibility mapping (QSM) with magnitude-, phase-, and R2\*-imaging at ultra-high magnetic field strength. *Neuroimage* 65, 299–314.
- Forstmann, B.U., Anwender, A., Schafer, A., Neumann, J., Brown, S., Wagenmakers, E.J., Bogacz, R., Turner, R., 2010. Cortico-striatal connections predict control over speed and accuracy in perceptual decision making. *Proc. Natl. Acad. Sci. U. S. A.* 107, 15916–15920.
- Forstmann, B.U., Keuken, M.C., Jahfari, S., Bazin, P.L., Neumann, J., Schafer, A., Anwender, A., Turner, R., 2012. Cortico-subthalamic white matter tract strength predicts interindividual efficacy in stopping a motor response. *Neuroimage* 60, 370–375.
- Gradinaru, V., Mogri, M., Thompson, K.R., Henderson, J.M., Deisseroth, K., 2009. Optical deconstruction of Parkinsonian neural circuitry. *Science* 324, 354–359.
- Greenhouse, I., Gould, S., Houser, M., Hicks, G., Gross, J., Aron, A.R., 2011. Stimulation at dorsal and ventral electrode contacts targeted at the subthalamic nucleus has different effects on motor and emotion functions in Parkinson's disease. *Neuropsychologia* 49, 528–534.
- Haynes, W.I., Haber, S.N., 2013. The organization of prefrontal-subthalamic inputs in primates provides an anatomical substrate for both functional specificity and integration: implications for basal ganglia models and deep brain stimulation. *J. Neurosci.* 33, 4804–4814.
- Hedreen, J.C., 1999. Tyrosine hydroxylase-immunoreactive elements in the human globus pallidus and subthalamic nucleus. *J. Comp. Neurol.* 409, 400–410.
- Hurd, Y.L., Suzuki, M., Sedvall, G.C., 2001. D1 and D2 dopamine receptor mRNA expression in whole hemisphere sections of the human brain. *J. Chem. Neuroanat.* 22, 127–137.
- Joel, D., Weiner, I., 1997. The connections of the primate subthalamic nucleus: indirect pathways and the open-interconnected scheme of basal ganglia-thalamocortical circuitry. *Brain Res. Brain Res. Rev.* 23, 62–78.
- Keuken, M.C., Uyllings, H.B., Geyer, S., Schafer, A., Turner, R., Forstmann, B.U., 2012. Are there three subdivisions in the primate subthalamic nucleus? *Front. Neuroanat.* 6, 14.
- Keuken, M.C., Bazin, P.L., Schafer, A., Neumann, J., Turner, R., Forstmann, B.U., 2013. Ultra-high 7 T MRI of structural age-related changes of the subthalamic nucleus. *J. Neurosci.* 33, 4896–4900.
- Kultas-Ilinsky, K., Leontiev, V., Whiting, P.J., 1998. Expression of 10 GABA(A) receptor subunit messenger RNAs in the motor-related thalamic nuclei and basal ganglia of Macaca mulatta studied with in situ hybridization histochemistry. *Neuroscience* 85, 179–204.
- Kuwajima, M., Hall, R.A., Aiba, A., Smith, Y., 2004. Subcellular and subsynaptic localization of group I metabotropic glutamate receptors in the monkey subthalamic nucleus. *J. Comp. Neurol.* 474, 589–602.
- Lambert, C., Zrinzo, L., Nagy, Z., Lutti, A., Hariz, M., Foltynie, T., Draganski, B., Ashburner, J., Frackowiak, R., 2012. Confirmation of functional zones within the human subthalamic nucleus: patterns of connectivity and sub-parcellation using diffusion weighted imaging. *Neuroimage* 60, 83–94.
- Langkammer, C., Schweser, F., Krebs, N., Deistung, A., Goessler, W., Scheurer, E., Sommer, K., Reishofer, G., Yen, K., Fazekas, F., Ropele, S., Reichenbach, J.R., 2012. Quantitative susceptibility mapping (QSM) as a means to measure brain iron? A post mortem validation study. *Neuroimage* 62, 1593–1599.
- Lee, Y., Dawson, V.L., Dawson, T.M., 2012. Animal models of Parkinson's disease: vertebrate genetics. *Cold Spring Harb. Perspect. Med.* 2.
- Levesque, J.C., Parent, A., 2005. GABAergic interneurons in human subthalamic nucleus. *Mov. Disord.* 20, 574–584.
- Mallet, L., Schupbach, M., N'Diaye, K., Remy, P., Bardinet, E., Czernecki, V., Welter, M.L., Pelissolo, A., Ruberg, M., Agid, Y., Yelnik, J., 2007. Stimulation of subterritories of the

- subthalamic nucleus reveals its role in the integration of the emotional and motor aspects of behavior. *Proc. Natl. Acad. Sci. U. S. A.* 104, 10661–10666.
- Massey, L.A., Miranda, M.A., Zrinzo, L., Al-Helli, O., Parkes, H.G., Thornton, J.S., So, P.W., White, M.J., Mancini, L., Strand, C., Holton, J.L., Hariz, M.I., Lees, A.J., Revesz, T., Yousry, T.A., 2012. High resolution MR anatomy of the subthalamic nucleus: imaging at 9.4 T with histological validation. *Neuroimage* 59, 2035–2044.
- Mori, S., Takino, T., Yamada, H., Sano, Y., 1985. Immunohistochemical demonstration of serotonin nerve fibers in the subthalamic nucleus of the rat, cat and monkey. *Neurosci. Lett.* 62, 305–309.
- Mundinger, F., 1965. Stereotaxic interventions on the zona incerta area for treatment of extrapyramidal motor disturbances and their results. *Confinia Neurologica* 26, 222–230.
- Navailles, S., Benazzouz, A., Bioulac, B., Gross, C., De Deurwaerdere, P., 2010. High-frequency stimulation of the subthalamic nucleus and L-3,4-dihydroxyphenylalanine inhibit in vivo serotonin release in the prefrontal cortex and hippocampus in a rat model of Parkinson's disease. *J. Neurosci.* 30, 2356–2364.
- Okun, M.S., Tagliati, M., Pourfar, M., Fernandez, H.H., Rodriguez, R.L., Alterman, R.L., Foote, K.D., 2005. Management of referred deep brain stimulation failures: a retrospective analysis from 2 movement disorders centers. *Arch. Neurol.* 62, 1250–1255.
- Parent, A., 1990. Extrinsic connections of the basal ganglia. *Trends Neurosci.* 13, 254–258.
- Parent, A., Hazrati, L.N., 1995a. Functional anatomy of the basal ganglia. I. The cortico-basal ganglia-thalamo-cortical loop. *Brain Res. Brain Res. Rev.* 20, 91–127.
- Parent, A., Hazrati, L.N., 1995b. Functional anatomy of the basal ganglia. II. The place of subthalamic nucleus and external pallidum in basal ganglia circuitry. *Brain Res. Brain Res. Rev.* 20, 128–154.
- Parent, A., Fortin, M., Cote, P.Y., Cicchetti, F., 1996. Calcium-binding proteins in primate basal ganglia. *Neurosci. Res.* 25, 309–334.
- Parent, M., Wallman, M.J., Gagnon, D., Parent, A., 2011. Serotonin innervation of basal ganglia in monkeys and humans. *J. Chem. Neuroanat.* 41, 256–265.
- Plaha, P., Ben-Shlomo, Y., Patel, N.K., Gill, S.S., 2006. Stimulation of the caudal zona incerta is superior to stimulation of the subthalamic nucleus in improving contralateral Parkinsonism. *Brain* 129, 1732–1747.
- Rafols, J.A., Fox, C.A., 1976. The neurons in the primate subthalamic nucleus: a Golgi and electron microscopic study. *J. Comp. Neurol.* 168, 75–111.
- Rughani, A.I., Hodaie, M., Lozano, A.M., 2013. Acute complications of movement disorders surgery: effects of age and comorbidities. *Mov. Disord.* 1661–1667.
- Schweser, F., Deistung, A., Lehr, B.W., Reichenbach, J.R., 2011. Quantitative imaging of intrinsic magnetic tissue properties using MRI signal phase: an approach to in vivo brain iron metabolism? *Neuroimage* 54, 2789–2807.
- Smith, Y., Hazrati, L.N., Parent, A., 1990. Efferent projections of the subthalamic nucleus in the squirrel monkey as studied by the PHA-L anterograde tracing method. *J. Comp. Neurol.* 294, 306–323.
- Temel, Y., Blokland, A., Steinbusch, H.W., Visser-Vandewalle, V., 2005. The functional role of the subthalamic nucleus in cognitive and limbic circuits. *Prog. Neurobiol.* 76, 393–413.
- Temel, Y., Boothman, L.J., Blokland, A., Magill, P.J., Steinbusch, H.W., Visser-Vandewalle, V., Sharp, T., 2007. Inhibition of 5-HT neuron activity and induction of depressive-like behavior by high-frequency stimulation of the subthalamic nucleus. *Proc. Natl. Acad. Sci. U. S. A.* 104, 17087–17092.
- Zaidel, A., Spivak, A., Grieb, B., Bergman, H., Israel, Z., 2010. Subthalamic span of beta oscillations predicts deep brain stimulation efficacy for patients with Parkinson's disease. *Brain* 133, 2007–2021.